



Effect of potassium and micronutrient fertilization on the activity of catalase and yield of wheat grown in saline conditions

Amal Al-Temimi¹; Saadi Al-Ghrairi^{1*}; Fadhil Al-Ghrairi¹; Ibrahim Razaq¹

1, Soil and Water Research Center, Directory of Agricultural Research, Ministry of Science and Technology, Baghdad, Iraq

Abstract

E-mail:
sadialgreary@yahoo.com

Received: 18/07/2020
Acceptance: 13/08/2020
Available Online: 20/08/2020
Published: 01/10/2020

Keywords: Wheat, Yield, Fertilization, Salinity, Catalase

This work was conducted as a pot experiment, to evaluate the effect of adding K, Fe, Zn, and Mn with conventional fertilizers on K/Na, Ca/Na and Mg/Na ratios and Catalase enzyme activity in leaves and on growth and yield of wheat (*Triticum aestivum L.*) under salinity conditions. K was added solely with conventional fertilizer dose (F1). In the second treatment, K, Fe, Zn, and Mn were sprayed on the plant as a foliar application before heading stage (F2). In the third treatment, K, Fe, Zn, and Mn were applied to the soil with conventional fertilizer dose (F3). Control treatment (F0), at which only conventional fertilizer dose was applied, for comparison. The field was irrigated with saline groundwater of 4.30 dS m⁻¹. Results showed that leaves K/Na, Ca/Na, and Mg/Na ratios as well as dry matter (F2) and (F3) treatments. Grain yield was increased by 29.60%, 31.52%, and 27.46% under F1, F2, and F3 treatments compared to that of control, respectively. Catalase activity under (F1) treatment was three times over that of control. The results suggested the importance of K and micronutrient fertilization when wheat is grown in salt-affected soil or irrigated with moderately saline water.

1. Introduction

The salinity of soil and irrigation water is one of the major stresses that limit plant growth and productivity in arid and semi-arid regions. Salinity usually induces nutritional disorders in plants which consequently affects plant biomass in addition to yield quantity and quality. It has been reported that these disorders may be the result of salinity direct effects on nutrient availability, competitive uptake, transport, or partitioning within the plant [1-4]. Salinity also affects plant growth since the high salt concentration in the soil solution interferes with the balanced absorption of the essential nutritional ions [5]. Furthermore, salinity causes imbalanced K⁺, Na⁺, Ca⁺², and Cl⁻ uptake in plants that alter the plant metabolism by affecting osmotic potential, enzymatic activities, membrane permeability, and electrochemical potential [6][7]. Therefore, improving the nutritional status of plants under salt stress conditions is of great importance for maintaining crop productivity.

Many studies reported that enhancement of plants nutrition status with mineral nutrients such as N, K, Ca, Mg, Fe, Zn, Mn, and Cu leads to increase crops tolerance to salinity due to the common physiological functions of these elements in plant cells such as the maintenance of photosynthesis activities and utilization of light energy in CO₂ fixation [4][8-16]

The salient effects of salinity in the plant are the impairment of the electron transport chain in the cell at high saline concentrations leading to the generation of reactive oxygen species (ROS). They are also called free radicals such as O₂⁻, H₂O₂, and OH. These powerful oxidizing agents quickly attack biological molecules such as DNA, lipids, and proteins, leading to severe disruption of cellular metabolic processes and damage to cell membranes [3][4]. To

prevent such degradation or damage, plant cells are equipped with anti-oxidant defense systems consisted of antioxidants such as non-enzymatic (carotenoids, glutathione, ascorbate & tocopherol) and antioxidant enzymes such as catalase (CAT), peroxidase (POD), and superoxide dismutase (SOD) which protects plant cells from oxidizing actions [17][18].

Catalase [$\text{H}_2\text{O}_2:\text{H}_2\text{O}_2$ oxidoreductase, EC (1.11.1.6)] is a tetrameric enzyme that follows the ping-pong mechanism to remove the harmful effects of H_2O_2 by removing the toxin complexes in the peroxide reactions and removing the electrons that form O_2^- . Usually, it is formed when the plant is subjected to bio-stress. In such circumstances, oxidative stress occurs due to the production and assembly of ROS such as O_2^- hydrogen peroxide H_2O_2 superoxide O_2^- hydroxide radical OH^- singlet oxygen $\text{O}^{\cdot-}$ [19][20].

Several studies have indicated the critical role of potassium ions and micronutrients Fe, Zn, and Mn in reducing the oxidative stress of the plant by stimulating or controlling the activity of antioxidant enzymes to reduce ROS production and also maintaining the electronic transmission chain in the process of photosynthesis [2][4][20-22]. Thus, improving the nutrient status of potassium and micro-elements in plants exposed to salt stress is essential to reduce oxidative stress that causes cell damage, at least by reducing ROS formation during photosynthesis [4]. Therefore, the objective of this work was to evaluate the effect of potassium and micronutrients in increasing yield and stimulating the activity of catalase enzyme (CAT) in wheat leaves grown under salt stress conditions.

2. Materials and Methods

2.1. Experiment and experimental design

A Pot experiment was carried out in the greenhouse of Soils & Water Resources Center, Directorate of Agricultural Research, Ministry of Science and Technology, Al-Zafaraniya, Baghdad, Iraq, to study the effect of potassium, iron, zinc and manganese fertilizers in the distribution ratios of potassium, calcium, magnesium to sodium and Catalase (CAT) activity in wheat (*Triticum aestivum* L.) (Tamose-2 variety) leaves when exposed to salt stress. Salt-affected soil with clay texture was collected from Al-Tuwaitha Research Station, 30 km southeast of Baghdad, classified as Typical Torrifluvent. Soil samples (0-30 cm) depth was air-dried, ground, and passed through a 2mm sieve mesh, thoroughly mixed and packed in plastic 6kg pots. The physicochemical properties of the soil before planting are presented in (Table 1). Minerals analysis was carried out according to the standard methods described by [23][24] using flame photometer, Spectrophotometer, and Atomic Absorption Spectrophotometer instruments.

Ten seeds of 'Tamose2' wheat variety were sown in each pot. Seedlings were thinned out to six per pot 15 days after germination.

The recommended fertilizers of nitrogen and phosphorus were added to control (F_0) and all other treatments in the forms of urea for N and triple superphosphate for P at the rates 100 and 30 mg kg⁻¹ soil, respectively. The fertilizers of potassium, iron, zinc, and manganese were added in the form K_2SO_4 , $\text{FeSO}_4 \cdot \text{H}_2\text{O}$, $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, and $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, respectively. The study included the following treatments:

F_0 : control

F_1 : Soil fertilization with potassium at 100 mg Kg⁻¹ soil in the form of K_2SO_4 (F_1).

F_2 : Foliar fertilizer of potassium, iron, zinc, and manganese in sulfate form were added with concentrations of 2500, 80, 60 and 40 mg l⁻¹ (2000, 64, 48 and 32 g ha⁻¹),

F_3 : Soil fertilization with potassium, iron, zinc, and manganese of 100, 20, 15, and 10 mg Kg⁻¹ soil respectively.

All phosphorus was added at sowing, while nitrogen was split into two equal doses at tillering and booting stages. Potassium, Iron, zinc, and manganese soil and foliar application were added at the booting stage. The treatments were

Table 1. Relevant chemical and physical properties of soil

Analysis	Value										
$\text{EC}_{(1:1)} \text{ dS.m}^{-1}$	7.24										
pH	7.28										
Soluble ions mmol.kg ⁻¹ soil	<table> <tr> <td>Na</td><td>42.80</td></tr> <tr> <td>Ca</td><td>7.00</td></tr> <tr> <td>Mg</td><td>8.10</td></tr> </table>	Na	42.80	Ca	7.00	Mg	8.10				
Na	42.80										
Ca	7.00										
Mg	8.10										
SAR	11.06										
$\text{CaCO}_3 \text{ gm.Kg}^{-1}$ soil	238										
CEC cm mol.Kg ⁻¹ soil	24.2										
N	48.30										
P	11.90										
Available elements mg.Kg ⁻¹ soil	<table> <tr> <td>K</td><td>132</td></tr> <tr> <td>Fe</td><td>1.43</td></tr> <tr> <td>Zn</td><td>0.38</td></tr> <tr> <td>Mn</td><td>0.34</td></tr> <tr> <td>Cu</td><td>0.88</td></tr> </table>	K	132	Fe	1.43	Zn	0.38	Mn	0.34	Cu	0.88
K	132										
Fe	1.43										
Zn	0.38										
Mn	0.34										
Cu	0.88										
Texture	Clay										
Field capacity (%)	34										

arranged in a completely randomized design (R.C.B.D.) in three replicates. All treatments were irrigated with brackish well water (Table 2) until harvest. The plants were harvested at maturity stage. Total dry matter and grains yield per pot were recorded.

2.2. Plant analysis

Random plant samples consisting of five flag leaves (blade & sheath) were collected from the main stem of a plant from each pot in the flowering stage, washed with normal water, and then distilled water to remove dust [25]. The leaves were cut into small pieces and dried at 65 °C for 48 hours and ground with an electric mill. 0.2 g of crushed dry leaves was digested with sulfuric and perchloric acids [26] and analyzed chemically for K, Na, Ca, and Mg elements according to the standard methods as in [24] using Atomic Absorption Spectrophotometer (400 P Analytica Jena, Germany).

2.3. Catalase activity assay

Samples of wheat leaves (5.0 g each) were collected at flowering stage. Leaf samples were homogenized in a potassium phosphate buffer solution (0.1M), pH=7.8 (1:2 w/v) using a pre-chilled mortar. The extract homogenate was transferred to centrifuge tubes and centrifuged at 12000g for 30 minutes and the supernatant was transferred to a new tube. The enzyme activity of the catalase (CAT: EC 1.11.1.6) was assayed using the method of [27]. 0.2 mL of the supernatant was incubated in 1 mL of substrate (65 µmol per mL H₂O₂ in 60 mmol/L sodium-potassium phosphate buffer, pH 7.4) at 37 °C for 1 min. The enzyme activity was suspended by adding 1 mL of 32.4 mM ammonium molybdate. The yellow absorption value of the molybdate complex and hydrogen peroxide was measured at 405 nm using a spectrophotometer (Specord205 Analytica Jena Germany). Catalase enzyme activity was calculated according to the following equation:

$$C.A (KU/l) = (S - B1/B2 - B3) * 271$$

C.A: Catalase activity (KU/l)

S: Sample reading.

B1: Blank 1 reading contained 1.0 ml substrate, 1.0 mL molybdate and 0.2 mL sample.

B2: Blank 2 reading contained 1.0 ml substrate, 1.0 ml molybdate, and 0.2 ml of 60 mmol/L sodium-potassium phosphate buffer, pH 7.4.

B3: Blank 3 reading contained 1.0 ml of 60 mmol/L sodium-potassium phosphate buffer pH 7.4, 1.0 mL molybdate and 0.2 ml of 60 mmol/L sodium-potassium phosphate buffer pH 7.4.

2.4. Statistical analysis

Data were analyzed statistically using Genstat software package and the means were compared using least significant difference (LSD 0.05).

3. Results and Discussion

3.1. Element concentrations ratio in wheat leaves

The effect of fertilization with potassium and microelements on K/Na, Ca/Na and Mg/Na ratio in leaves of wheat during the flowering stage is given in (Table3). Results showed that the concentration ratios in the wheat leaves were significantly affected by the application of K, Fe, Zn, and Mn. This application induced an increase in K/Na, Ca/Na, and Mg/Na ratios in leaves. The increase was more pronounced in K/Na ratio which was 2.44, 2.73, 3.04, and 3.60 under F0, F1, F2, and F3 treatment, respectively. The highest K/Na ratios were obtained under (F2) and (F3) treatments. This increase in K/Na ratio may be attributed to trace elements' role in increasing the capability of the root system for

Table 2. Properties of well water used in irrigation

EC	pH	Soluble Ions mmol. l ⁻¹					Sodium adsorption ratio (SAR)
		SO ₄ ²⁻	Cl ⁻¹	Mg ⁺²	Ca ⁺²	Na ⁺	
4.30	7.32	8.08	23.13	6.02	5.35	21.22	6.30

selectivity uptake of K⁺ ion and the limitation of Na ion uptake in the leaves [28][29]. This indicates that the addition of potassium and trace elements with regular fertilizer dose can increase salt tolerance of wheat by reducing Na uptake. [1][30] reported that the plant which is capable of maintaining a high ratio of K/Na has high salinity tolerance.

Ca/Na and Mg/Na ratios were relatively equal. Ca/Na ratio was 1.66 under foliar fertilization treatment (F2) which is significantly higher than the other three treatments. Results also showed no significant differences among Ca/Na ratios under F1 and F0 treatments. Additionally, Mg/Na ratio was significantly higher in (F2) treatment (1.48) when compared to the three other treatments.

These results are consistent with previous results that ground or foliar fertilization with major and minor nutrients has resulted in improved absorption of potassium, calcium, and magnesium elements [10-16][21] resulting in an increase in plants tolerance to saline stress conditions as confirmed by [31].

3.2. Dry matter and grains yield

Effect of the selected fertilization treatments on dry matter and grain yield (g pot⁻¹) of wheat is given in (Table 4). Results indicated that there was a significant effect of potassium and trace elements application on the total dry matter. The highest dry matter was recorded under F2 treatment (34.2 g pot⁻¹). On the other hand, the lowest dry matter value was (29.40 g pot⁻¹) was obtained under treatment F0 treatment. The recorded increases in the dry matter over that of control (F0) were 12.22, 16.20, and 13.24% under F1, F2, and F3 fertilization treatments, respectively.

The grain yield (Table 4) was significantly increased under F1, F2, and F3 treatments in comparison to control, and the highest grain yield was 6.80 g pot⁻¹ under F2 treatment (foliar application). The increase in the yield over that of control was 29.60, 31.52, and 27.46% under F1, F2, and F3 treatments, respectively. That indicates the importance of foliar application of K and trace elements in alleviating the adverse effect of salt stress on growth and grain production of wheat which is in correspondence with previous studies [4][10][11][14][16][21][32].

3.3. Catalase enzyme activity (CAT)

Higher CAT activity was observed under the soil application of K (F1) compared to other treatments (181.98 KU.l⁻¹) which was three folds higher than that of the control treatment (F0). On the other hand, enzyme activity under conventional fertilizer dose was 58.03 KU.l⁻¹,

which is significantly lower than other treatments (Table 5). The differences between F2 (foliar application) and F3 can be explained by the fact that concentration of Mn, Zn, Fe, Ca, and K nutrients in wheat plants at F3 treatment were not enough to increase the enzyme's activity to counteract salt stress. This may be attributed to the precipitate part of the added micronutrients due to higher pH and carbonate content in the used clay soil [33]. The increments in CAT activity helped plant in scavenging H₂O₂ which is generated in normal or abnormal conditions. Additionally, CAT activity maintained the ascorbate pool and activated the defense system of plant cells which in turn led to an increase in wheat plant tolerance and maintained the best growth under saline conditions [2][34]. These results confirm that this enzyme is more effective in wheat when K, Fe, Zn, and Mn were added either to the soil or foliar application. This is due to the fact that K and micronutrients are necessary for the synthesis of proteins and the effectiveness of enzymes as it is a carrier element and acts to change the shape and configuration of the active site in the enzyme molecule to obtain better binding between the substrate and the enzyme [35-37].

Table 3. Effect of fertilization with potassium and micronutrients on elements ratios in wheat leaves

The ratio	Treatments				LSD(0.05)
	F ₀	F ₁	F ₂	F ₃	
K/Na	2.44	2.73	3.04	3.60	0.06
Ca/Na	1.09	1.26	1.66	1.52	0.30
Mg/Na	0.80	1.11	1.48	1.33	0.09

Table 4. Effect of fertilization with potassium and micronutrients on the dry matter and grains yield (g pot⁻¹) of wheat

The yield	Treatments				LSD(0.05)
	F ₀	F ₁	F ₂	F ₃	
Dry matter (g pot ⁻¹)	29.39	32.98	34.15	33.28	0.09
Grains (g pot ⁻¹)	5.17	6.70	6.80	6.59	0.11

Table 5. Effect of fertilization with potassium and micronutrients in catalase enzyme activity CAT (KU l⁻¹) in wheat leaves

Fertilization treatment	F ₀	F ₁	F ₂	F ₃
Enzyme activity (KU l ⁻¹)	58.03	181.98	179.53	134.78
LSD(0.05)		1.65		

Conflict of interest statement

The authors declared no conflict of interest.

Funding statement

The authors declared that no funding was received in relation to this manuscript.

Data availability statement

The authors declared that all related data are included in the article.

References

1. Abbasi GH, Akhtar J, Ahmad R, Jamil M, Anwar-ul-Haq M, Ali S, Ijaz M. Potassium application mitigates salt stress differentially at different growth stages in tolerant and sensitive maize hybrids. *Plant Growth Regul.* 2015;76(1):111-25. [DOI](#)
2. Al-Samerria IK, Al-Ghrairi SM, Rahi HAS. Induction of antioxidant enzymes in wheat (*Triticum spp.*) grown under salt stress. *Baghdad J. Sci.* 2013;10:832-43.
3. Hernandez JA, Jiménez A, Mullineaux P, Sevilia F. Tolerance of pea (*Pisum sativum L.*) to long-term salt stress is associated with induction of antioxidant defences. *Plant Cell Environ.* 2000;23(8):853-62. [DOI](#)
4. Cakmak I. The role of potassium in alleviating detrimental effects of abiotic stresses in plants. *J Plant Nutr Soil Sc.* 2005;168(4):521-30. [DOI](#)
5. Tester M, Davenport R. Na tolerance and Na transportation in higher plants. *Ann Bot.* 2003;91(5):503-27.
6. Grattan SR, Grieve CM. Salinity-mineral nutrient relations in horticultural crops. *Sci Hortic-Amsterdam.* 1998;78(1-4):127-57. [DOI](#)
7. Khan MA, Ashraf MY, Mujtaba SM, Shirazi MU, Khan MA, Shereen A, Mumtaz S, Siddiqui MA, Kaleri GM. Evaluation of high yielding canola type *Brassica* genotypes/mutants for drought tolerance using physiological indices as screening tool. *Pak. J. Bot.* 2010 Dec 1;42(6):3807-16.
8. Abou-El-Nour EZ, Aly EM, El-Fouly MM, Salama ZA. Chelated Fe and Zn foliar spray improve the tolerance of kidney bean (var. nebraska) plants in salinized media. *Biosci. Res.* 2017 Jul 1;14(3):525-31.
9. El-Fouly MM, Mobarak ZM, Salama ZA. Micronutrients (Fe, Mn, Zn) foliar spray for increasing salinity tolerance in wheat *Triticum aestivum L.* *Afr. J. Plant Sci.* 2011;5(5):314-22. [DOI](#)
10. Al-Ghrairi SM. Alleviation of the adverse effect of salt stress on growth and yield of wheat by using foliar application. (Ph.D. thesis, Baghdad Univirstry). 2011.
11. Al-Ghrairi SM, Razzaq IB, Al-Hasani AMM, Khudhier SA, Jasim BA, Mousa RA. Effect Of Some Fertilizer Combinations On Growth And Yield Of Wheat Grown In Salt -Affected Soil. The 10th Scientific Conference for Agricultural research. Baghdad, Iraq. 2017.
12. Al-Magrebi NM. Effect potassium and phosphorus fertilizer on growth and yield of *Sorghum bicolor (L.) moench* irrigated with deferent irrigation water salinity. (Ph.D. thesis, Baghdad University). 2004.
13. Genc Y, McDonald GK, Tester M. Reassessment of tissue Na⁺ concentration as a criterion for salinity tolerance in bread wheat. *Plant Cell Environ.* 2007;30(11):1486-98. [DOI](#)

14. Manal FM, Thalooth AT, Khalifa RK. Effect of foliar spraying with uniconazole and micronutrients on yield and nutrients uptake of wheat plants grown under saline condition. Am. J. Sci. 2010;6(8):398-404.
15. Zaman B, Niazi BH, Athar M, Ahmad M. Response of wheat plants to sodium and calcium ion interaction under saline environment. Int. J. Environ. Sci. 2005;2(1):7-12.
16. Abd El-Hady BA. Effect of zinc application on growth and nutrient uptake of barley plant irrigated with saline water. Res. J. Appl. Sci. 2007;3(6):431-6.
17. Shalata A, Tal M. The effect of salt stress on lipid peroxidation and antioxidants in the leaf of the cultivated tomato and its wild salt-tolerant relative *Lycopersicon pennellii*. Physiol. Plant. 1998;104(2):169-74. [DOI](#)
18. Meloni DA, Oliva MA, Martinez CA, Cambraia J. Photosynthesis and activity of superoxide dismutase, peroxidase and glutathione reductase in cotton under salt stress. Environ. Exp. Bot. 2003;49(1):69-76. [DOI](#)
19. Kuk YI, Shin JS, Burgos NR, Hwang TE, Han O, Cho BH, Jung S, Guh JO. Antioxidative enzymes offer protection from chilling damage in rice plants. Crop Sci. 2003;43(6):2109-17. [DOI](#)
20. Cui Y, Zhao N. Oxidative stress and change in plant metabolism of maize (*Zea mays L.*) growing in contaminated soil with elemental sulfur and toxic effect of zinc. Plant Soil Environ. 2011;57(1):34-9. [DOI](#)
21. Al-Ghrairi SM, Al-Samerria IK, Rahi HS. Effect of salt stress and foliar application of some nutrients on K and Ca accumulation , distribution , and its relation on peroxidase enzyme activity in leaves of wheat. Iraq J. Sci. Tech. 2014;5:20-9.
22. Moran JF, James EK, Rubio MC, Sarath G, Klucas RV, Becana M. Functional characterization and expression of a cytosolic iron-superoxide dismutase from cowpea root nodules. Plant Physiol. 2003;133(2):773-82. [DOI](#)
23. Richards LA. Diagnosis and improvement of saline and alkali soils. Handbook No. 60. US Department of Agriculture, Washington, DC. 1954.
24. Page AL, Miller RH, Keeney DR. Chemical and microbiological properties. Methods of soil analysis. 1982;2.
25. Loop EA, Finck A. Total iron as a useful index of the Fe-status of crops. J. Plant Nutr. 1984;7(1-5):69-79. [DOI](#)
26. Gresser MS, Parson GW. Sulfuric. pere chloric acid digestion of plant material for the determination nitrogen , phosphorus , potassium , calcium and Mg . Anal. Chim. Acta. 1979;109:431-6.
27. Goth L. A simple method for determination of serum catalase activity and revision of reference range. Clin. Chim. Acta. 1991;196(2-3):143-51. [DOI](#)
28. Carvajal M, Martinez V, Cerda A. Influence of magnesium and salinity on tomato plants grown in hydroponic culture. J. Plant Nutr. 1999;22(1):177-90. [DOI](#)
29. Zheng Y, Jia A, Ning T, Xu J, Li Z, Jiang G. Potassium nitrate application alleviates sodium chloride stress in winter wheat cultivars differing in salt tolerance. J. Plant Physiol. 2008;165(14):1455-65. [DOI](#)
30. Khan MA, Shirazi MU, Khan MA, Mujtaba SM, Islam E, Mumtaz S, Shereen A, Ansari RU, Ashraf MY. Role of proline, K/Na ratio and chlorophyll content in salt tolerance of wheat (*Triticum aestivum L.*). Pak. J. Bot. 2009 Apr 1;41(2):633-8.
31. Bortner CD, Cidlowski JA. Cell shrinkage and monovalent cation fluxes: role in apoptosis. Arch. Biochem. Biophys. 2007;462(2):176-88. [DOI](#)

32. Al-Samerria IK, Al-Ghrairi SM, Rahi HS. Effect of salt stress and foliar application of some nutrients and its relation on superoxidase enzyme activity and proline content in growth of wheat plant. *Iraq J. soil sci.* 2013;13:132-40.
33. Lindsay WL. Chemical equilibria in soils. John Wiley&Sons, NewYork. 1979.
34. Amal GA, Orabi S, Gomaa AM. Bio-organic farming of grain sorghum and its effect on growth, physiological and yield parameters and antioxidant enzymes activity. *Res. j. agric. biol. sci.* 2010;6(3):270-9.
35. Van Brunt JM, Sultenfuss JH. Better crops with plant food. Potassium: Functions of potassium. 1998;82(3):4-5.
36. Sangakkara UR, Frehner M, Nösberger J. Influence of soil moisture and fertilizer potassium on the vegetative growth of mungbean (*Vigna radiata* L. Wilczek) and cowpea (*Vigna unguiculata* L. Walp). *J Agron Crop Sci.* 2001;186(2):73-81. [DOI](#)
37. Mahmood K. Salinity tolerance in barley (*Hordeum vulgare* L.): effects of varying NaCl, K⁺/Na⁺ and NaHCO₃ levels on cultivars differing in tolerance. *Pak J Bot.* 2011;43(3):1651-4.